

Integral representation of some functions related to the Gamma function

Christian Berg

Abstract. We prove that the functions $\Phi(x) = [\Gamma(x+1)]^{1/x}(1+1/x)^x/x$ and $\log \Phi(x)$ are Stieltjes transforms.

Mathematics Subject Classification (2000). Primary 33B15; Secondary 26A48.

Keywords. Complete monotonicity, Gamma function.

1. Introduction and main results

In [11] the authors introduce a subclass of the completely monotonic functions which they call *logarithmically completely monotonic*, and the main result in [12] is that the function

$$\Phi(x) = \frac{[\Gamma(x+1)]^{1/x}}{x} \left(1 + \frac{1}{x}\right)^x \quad (1)$$

is logarithmically completely monotonic.

We characterize the class of logarithmically completely monotonic functions as the infinitely divisible completely monotonic functions studied by Horn in [14]. We prove that Stieltjes transforms (see (8) below) belong to this class and that Φ and $\log \Phi$ are both Stieltjes transforms. Each of these statements imply the result of [12]. The following explicit representations are obtained:

$$\log \Phi(x) = \int_0^\infty \frac{\varphi(s)}{s+x} ds, \quad x > 0, \quad (2)$$

where

$$\varphi(s) = \begin{cases} 1-s & \text{if } 0 \leq s < 1 \\ 1-n/s & \text{if } n \leq s < n+1, n=1,2,\dots \end{cases} \quad (3)$$

and

$$\Phi(x) = 1 + \int_0^\infty \frac{h(s)}{s+x} ds, \quad x > 0, \quad (4)$$

with

$$h(s) = \frac{1}{\pi} \frac{s^{s-1}}{|1-s|^s |\Gamma(1-s)|^{1/s}} \sin(\pi\varphi(s)), \quad s \geq 0. \quad (5)$$

Note that the density $\varphi(s)$ takes its values in the interval $[0, 1]$, and this is the clue to the fact that also Φ is a Stieltjes transform. The density h is continuous on $[0, \infty[$ with $h(0) = \exp(-\gamma)$, where γ is Euler's constant, and $h(n) = 0$ for $n \in \mathbb{N}$.

Recall that a function $f :]0, \infty[\rightarrow \mathbb{R}$ is said to be completely monotonic, if f has derivatives of all orders and satisfies

$$(-1)^n f^{(n)}(x) \geq 0 \quad \text{for all } x > 0 \quad \text{and } n = 0, 1, 2, \dots$$

Bernstein's Theorem, cf. [15, p. 161], states that f is completely monotonic if and only if

$$f(x) = \int_0^\infty e^{-xs} d\mu(s), \quad (6)$$

where μ is a nonnegative measure on $[0, \infty)$ such that the integral converges for all $x > 0$. The set of completely monotonic functions is denoted $\mathcal{C}$.

In [12] the authors call a function $f :]0, \infty[\rightarrow]0, \infty[$ *logarithmically completely monotonic* if it is C^∞ and

$$(-1)^k [\log f(x)]^{(k)} \geq 0, \text{ for } k = 1, 2, \dots \quad (7)$$

If we denote the class of logarithmically completely monotonic functions by $\mathcal{L}$, we have $f \in \mathcal{L}$ if and only if f is a positive C^∞ -function such that $-(\log f)' \in \mathcal{C}$.

The functions of class $\mathcal{L}$ have been implicitly studied in [3], and Lemma 2.4(ii) in that paper can be stated as the inclusion $\mathcal{L} \subset \mathcal{C}$, a fact also established in [11].

The class $\mathcal{L}$ can be characterized in the following way, established by Horn[14, Theorem 4.4]:

Theorem 1.1. *For a function $f :]0, \infty[\rightarrow]0, \infty[$ the following are equivalent:*

- (i) $f \in \mathcal{L}$
- (ii) $f^\alpha \in \mathcal{C}$ for all $\alpha > 0$
- (iii) $\sqrt[n]{f} \in \mathcal{C}$ for all $n = 1, 2, \dots$

Another way of expressing the conditions of Theorem 1.1 is that the functions in $\mathcal{L}$ are those completely monotonic functions for which the representing measure μ in (6) is infinitely divisible in the convolution sense: For each $n \in \mathbb{N}$ there exists a positive measure ν on $[0, \infty[$ with n 'th convolution power equal to μ , viz. $\nu^{*n} = \mu$. By condition (ii) there exists a convolution semigroup $(\mu_\alpha)_{\alpha > 0}$ of positive measures such that the Laplace transform of μ_α is f^α . Note that the convolution of any two positive measures on $[0, \infty[$ is well-defined and we have $\mu_\alpha * \mu_\beta = \mu_{\alpha+\beta}$.

In the special case of $f(0+) = 1$ this is very classical: This is the description of infinitely divisible distributions in probability. Since there are probabilities which are not infinitely divisible we have $\mathcal{C} \setminus \mathcal{L} \neq \emptyset$.

In various papers complete monotonicity for special functions has been established by proving the stronger statement that the function is a Stieltjes transform, i.e. is of the form

$$f(x) = a + \int_0^\infty \frac{d\mu(s)}{s+x}, \quad (8)$$

where $a \geq 0$ and μ is a nonnegative measure on $[0, \infty[$ satisfying

$$\int_0^\infty \frac{1}{1+s} d\mu(s) < \infty.$$

See [2],[3],[5],[6],[8],[9].

The set of Stieltjes transforms will be denoted $\mathcal{S}$. We clearly have $\mathcal{S} \subset \mathcal{C}$. For more information about this class see [7].

Theorem 1.2. $\mathcal{S} \setminus \{0\} \subset \mathcal{L}$.

Theorem 1.3. *The functions*

$$\Phi(x) = \frac{[\Gamma(x+1)]^{1/x}}{x} \left(1 + \frac{1}{x}\right)^x$$

and

$$\log \Phi(x) = \frac{\log \Gamma(x+1)}{x} - \log x + x \log \left(1 + \frac{1}{x}\right)$$

are Stieltjes transforms with the representations (4) and (2).

Remark 1.4. The class $\mathcal{S}$ has the following stability properties: If $f \in \mathcal{S}, f \neq 0$ then $1/f(1/x)$ and $1/(xf(x))$ are again Stieltjes transforms, cf. [5]. Therefore the following functions belong to $\mathcal{S}$:

$$\frac{1}{[\Gamma(1+x)]^{1/x}(1+1/x)^x}, \quad [\Gamma(1+1/x)]^x(1+x)^{1/x}, \quad \frac{1}{x[\Gamma(1+1/x)]^x(1+x)^{1/x}}.$$

It was proved in [3] that $[\Gamma(1+1/x)]^x \in \mathcal{S}$, so also the following functions are Stieltjes transforms:

$$\frac{[\Gamma(1+x)]^{1/x}}{x}, \quad \frac{1}{[\Gamma(1+x)]^{1/x}}, \quad \frac{1}{x[\Gamma(1+1/x)]^x}.$$

In [2] it was proved that $(1+1/x)^{-x} \in \mathcal{S}$. Therefore the function Φ given by (1) is a quotient of known Stieltjes transforms, but this does not imply that the function itself is a Stieltjes transform.

2. Proofs

For completeness we include a proof of Theorem 1.1.

Proof. "(i) $\Rightarrow$ (ii)". Since $f \in \mathcal{L}$ implies $f^\alpha \in \mathcal{L}$ for all $\alpha > 0$, it is enough to prove the inclusion $\mathcal{L} \subset \mathcal{C}$. Although this is done in [3] and [11] we include the easy proof. By assumption $-(\log f)' = -f'/f \in \mathcal{C}$, so in particular $-f' \geq 0$. Assume now that $(-1)^k f^{(k)} \geq 0$ for $k \leq n$. Then

$$\begin{aligned} (-1)^{n+1} f^{(n+1)} &= (-1)^n ((-\log f)' f)^{(n)} \\ &= \sum_{k=0}^n \binom{n}{k} (-1)^k ((-\log f)')^{(k)} (-1)^{n-k} f^{(n-k)} \geq 0, \end{aligned}$$

and (ii) follows by induction.

"(ii) $\Rightarrow$ (iii)" is obvious.

"(iii) $\Rightarrow$ (i)". If $f^{1/n} \in \mathcal{C}$ we have in particular $-(f^{1/n})' = (-1/n)f^{-1+1/n}f' \in \mathcal{C}$. Multiplying by n and letting $n \rightarrow \infty$ we see that the limit function $-f'/f$ belongs to $\mathcal{C}$, because $\mathcal{C}$ is closed under pointwise limits, cf. [7]. This establishes (i). $\square$

Proof of Theorem 1.2: Let $f \in \mathcal{S}$ be non-zero, and let $\alpha > 0$. By Theorem 1.1 it is enough to prove that $f^\alpha \in \mathcal{C}$. Writing $\alpha = n + a$ with $n = 0, 1, \dots$ and $0 \leq a < 1$ we have $f^\alpha = f^n f^a$, and using the stability of $\mathcal{C}$ under multiplication and that $f^a \in \mathcal{S}$, cf. [4], the assertion follows. $\square$

Proof of Theorem 1.3: Using the expression (3) for φ we find

$$\begin{aligned} \int_0^\infty \frac{\varphi(s)}{s+x} ds &= \int_0^1 \frac{1-s}{s+x} ds + \sum_{k=1}^\infty \int_k^{k+1} \frac{1-k/s}{s+x} ds \\ &= -1 + (x+1) \log \left(1 + \frac{1}{x} \right) \\ &\quad + \sum_{k=1}^\infty \left[\left(1 + \frac{k}{x} \right) \log \left(1 + \frac{1}{x+k} \right) - \frac{k}{x} \log \left(1 + \frac{1}{k} \right) \right]. \end{aligned}$$

Therefore,

$$\int_0^\infty \frac{\varphi(s)}{s+x} ds = \log \Phi(x)$$

if and only if

$$\log \Gamma(x+1) = x(\log(1+x) - 1) + \sum_{k=1}^{\infty} \left[(k+x) \log \left(1 + \frac{1}{x+k} \right) - k \log \left(1 + \frac{1}{k} \right) \right] \quad (9)$$

for $x \geq 0$. Both sides vanish for $x = 0$, and they have the same derivative $\psi(x+1)$, where ψ is the digamma function. This follows easily by the classical formula

$$\psi(x) = \log x + \sum_{k=0}^{\infty} \left[\log \left(1 + \frac{1}{x+k} \right) - \frac{1}{x+k} \right]$$

cf. [13, 8.362(2)]. This shows that $\log \Phi$ is a Stieltjes transform with the representation (2). In particular Φ is completely monotonic with the limit 1 at infinity.

To see that a function f is a Stieltjes transform we will use the characterization of these functions via complex analysis, see [1, p. 127] or [5]. It is necessary and sufficient that f has a holomorphic extension to the cut plane $\mathcal{A} = \mathbb{C} \setminus]-\infty, 0]$ and satisfies $\operatorname{Im} f(z) \leq 0$ for $\operatorname{Im} z > 0$ and $f(x) \geq 0$ for $x > 0$. For a Stieltjes transform f given by (8) we have $a = \lim_{x \rightarrow \infty} f(x)$, and the measure μ is the limit in the vague topology of

$$-\frac{1}{\pi} \operatorname{Im} f(-x+iy) dx$$

as $y \rightarrow 0^+$.

We clearly have $\Phi(x) > 0$ for $x > 0$ and the holomorphic extension of Φ is given by $\Phi(z) = \exp(\log \Phi(z))$, where $\log \Phi(z)$ is the holomorphic extension obtained by the representation (2). This can also be described in the following way: For $z \in \mathcal{A}$ we let $\log \Gamma(z)$ be the unique holomorphic branch, which is real for $x > 0$, and we let Log denote the principal logarithm. Then

$$\frac{\log \Gamma(z+1)}{z} - \operatorname{Log} z + z \operatorname{Log} \left(1 + \frac{1}{z} \right) \quad (10)$$

is a holomorphic branch of $\log \Phi(z)$ in $\mathcal{A}$, and since it agrees with $\log \Phi(x)$ for $x > 0$, we have

$$\log \Phi(z) = \frac{\log \Gamma(z+1)}{z} - \operatorname{Log} z + z \operatorname{Log} \left(1 + \frac{1}{z} \right) = \int_0^{\infty} \frac{\varphi(s)}{s+z} ds, \quad z \in \mathcal{A}. \quad (11)$$

For $z = x+iy, y > 0$ we get

$$\operatorname{Im} \log \Phi(x+iy) = - \int_0^{\infty} \frac{\varphi(s)y}{(s+x)^2 + y^2} ds,$$

and since $0 \leq \varphi(s) \leq 1$ for $s \geq 0$ we get

$$\operatorname{Im} \log \Phi(x+iy) \in]-\pi, 0[,$$

hence

$$\operatorname{Im} \Phi(x+iy) = |\Phi(x+iy)| \sin(\operatorname{Im} \log \Phi(x+iy)) < 0,$$

which shows that Φ is a Stieltjes transform. For $x \geq 0, y \rightarrow 0^+$ we further get

$$-\frac{1}{\pi} \operatorname{Im} \Phi(-x+iy) \rightarrow h(x) := \frac{1}{\pi} |\Phi(-x)| \sin(\pi \varphi(x)), \quad (12)$$

which is a continuous nonnegative function on $[0, \infty[$. Therefore the convergence is uniform for x in compact subsets of $[0, \infty[$, so h is the density of the representing measure as a Stieltjes transform.

This shows the following integral representation of Φ :

$$\Phi(z) = 1 + \int_0^{\infty} \frac{h(s)}{s+z} ds, \quad z \in \mathcal{A} \quad (13)$$

where

$$h(s) = \frac{1}{\pi} \frac{s^{s-1}}{|1-s|^s |\Gamma(1-s)|^{1/s}} \sin(\pi \varphi(s)), \quad s \geq 0.$$

□

Remark 2.1. There is a close relationship between Stieltjes transforms and Pick functions. For the latter see [1] and [10]. It is possible to find the integral representation (2) of $\log \Phi(x)$ using integral representations of the three terms of (10). Here $\text{Log}(z)$ and $z \text{Log}(1 + 1/z)$ are Pick functions for which the integral representations are easily obtained. The integral representation of the Pick function $(\text{Log } \Gamma(z+1))/z$ was found in [9]. The author first found the density (3) in this way, but once φ is found the present direct approach seems easier.

References

- [1] N. I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis*. English translation, Oliver and Boyd, Edinburgh, 1965.
- [2] H. Alzer, C. Berg, *Some Classes of Completely Monotonic Functions*, Ann. Acad. Sci. Fenn. Math. **27**, (2002), 445–460.
- [3] H. Alzer, C. Berg, *Some Classes of Completely Monotonic Functions, II*. To appear in The Ramanujan Journal.
- [4] C. Berg, *The Stieltjes cone is logarithmically convex*, In: Complex Analysis Joensuu 1978. Lecture Notes in Mathematics **747**, 46–54. Springer, Berlin-Heidelberg-New York, 1979.
- [5] C. Berg, *Quelques remarques sur le cône de Stieltjes*, in: Séminaire de Théorie du Potentiel Paris, No. 5, Lecture Notes in Mathematics **814**, Springer, Berlin-Heidelberg-New York, 1980.
- [6] C. Berg, *A Pick function related to an inequality for the entropy function*, J. Inequal. Pure Appl. Math. **2**, Issue 2, Article 26 (2001).
- [7] C. Berg, G. Forst, *Potential theory on locally compact abelian groups*. Springer-Verlag, Berlin-Heidelberg-New York, 1975.
- [8] C. Berg, H.L. Pedersen, *A completely monotone function related to the Gamma function*, J. Comput. Appl. Math. **133**, (2001), 219–230.
- [9] C. Berg, H.L. Pedersen, *Pick functions related to the gamma function*, Rocky Mount. J. Math. **32** (2002), 507–525.
- [10] W.F. Donoghue, Jr., *Monotone Matrix Functions and Analytic Continuation*. Springer-Verlag, Berlin-Heidelberg-New York, 1974.
- [11] Feng Qi, Bai-Ni Guo, *Complete monotonicities of functions involving the Gamma and Digamma functions*, RGMIA Res. Rep. Coll. **7**, No. 1 (2004), Art. 6. Available online at <http://rgmia.vu.edu.au/v7n1.html>
- [12] Feng Qi, Bai-Ni Guo, and Chao-Ping Chen, *Some completely monotonic functions involving the Gamma and polygamma functions* RGMIA Res. Rep. Coll. **7**, No. 1 (2004), Art. 8. Available online at <http://rgmia.vu.edu.au/v7n1.html>
- [13] I.S. Gradshteyn, I.M. Ryzhik, *Table of Integrals, Series and Products*. Sixth Edition, Academic Press, San Diego, 2000.
- [14] R.A. Horn, *On infinitely divisible matrices, kernels and functions*, Z. Wahrscheinlichkeitstheorie und Verw. Geb. **8** (1967), 219–230.
- [15] D.V. Widder, *The Laplace Transform*. Princeton University Press, Princeton, 1941.

Christian Berg
 Department of Mathematics
 University of Copenhagen
 Universitetsparken 5
 DK-2100 Copenhagen
 Denmark
 e-mail: berg@math.ku.dk